

Conference Presentation

The effect of ultra high-diluted drugs on plant-nematode interaction

Natalia Rodiuc

Laboratório de Interação Molecular Planta-Praga, Embrapa Recursos Genéticos e Biotecnologia, PqEB, Av. W5 Norte final, Brasília/DF, 70770-900, Brazil

Institut National de la Recherche Agronomique, UMR 1355 ISA/Centre National de la Recherche Scientifique, UMR 7254 ISA/Université de Nice-Sophia Antipolis, UMR ISA, 400 route des Chappes, Sophia-Antipolis, France

Email: Natalia.Ostanin@sophia.inra.fr

Abstract

Among plant pathogens, sedentary endoparasitic nematodes are one of the most damaging pests in global agriculture. Within this group, root-knot nematodes (RKN) *Meloidogyne* spp. is one of the most specialized phytoparasitic nematodes according to the complexity of the induced feeding sites. These phytopathogenic worms are highly resistant due to the large physiological variability, therefore difficult to fight against. The traditional methods of control, such as the crop rotation and the use of resistant varieties by classical selection are not always effective, time consuming and costly. The modern methods including the application of genetically modified crops (GMO), synthetic pesticides and bionematicides could be dangerous for human health and ecologically hazardous (Thomason, 1987; Malatesta *et al.* 2002a,b, 2008a,b; Spiroux de Vendômois *et al.*, 2010; Seralini *et al.*, 2012). Therefore, alternative routes must be taken in order to obtain plants resistant to nematodes. The ultra high dilutions (HDs) approach is largely in charge to dissect the infective process. HDs method is eco-friendly, low cost and leaves no residue in the environment. The classical phytopathological methods combined with modern microscopy approaches allowed to characterize the effect of different HD drugs on compatible interaction between model plant *Arabidopsis thaliana* and root-knot nematodes *Meloidogyne incognita*.

The work is under consideration for publication elsewhere.

References

1. [Malatesta M, Caporaloni C, Gavaudan S, Rocchi MB, Serafini S, Tiberi C and Gazzanelli G \(2002a\). Ultrastructural morphometrical and immunocytochemical analyses of hepatocyte nuclei from mice fed on genetically modified soybean. *Cell Struct. Funct.* 27: 173-180.](#)
2. [Malatesta M, Caporaloni C, Rossi L, Battistelli S, Rocchi MB, Tonucci F and Gazzanelli G \(2002b\). Ultrastructural analysis of pancreatic acinar cells from mice fed on genetically modified soybean. *J. Anat.* 201\(5\): 409-415.](#)
3. [Malatesta M, Boraldi F, Annovi G, Baldelli B, Battistelli S, Biggiogera M and Quaglino D \(2008a\). A long-term study on female mice fed on a genetically modified soybean: effects on liver ageing. *Histochem. Cell Biol.* 130\(5\): 967-977.](#)



Cite as: Rodiuc N. The effect of ultra-high-diluted drugs on plant-nematode interaction. Proceedings of the XXIX GIRI Meeting; 2015 June 3 – 5; Verona (Italy). *Int J High Dilution Res.* 2015; 14(2): 9-10

-
4. [Malatesta M, Perdoni F, Santin G, Battistelli S, Muller S and Biggiogera M \(2008b\). Hepatoma tissue culture \(HTC\) cells as a model for investigating the effects of low concentrations of herbicide on cell structure and function. *Toxicol. In Vitro* 22\(8\): 1853-60.](#)
 5. [Seralini GE, Clair E, Mesnage R, Gress S, Defarge N, Malatesta M, Hennequin D and de Vendomois JS \(2012\). Long term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. *Food Chem Toxicol.* 2012; 50\(11\): 4221-31.](#)
 6. [Vendômois JS, Roullier F, Cellier D and Seralini GE \(2009\). A comparison of the effects of three GM corn varieties on mammalian health. *Int. J. Biol. Sci.* 2009;5\(7\):706-26.](#)
 7. Thomason IJ (1987). Challenges facing nematology: environmental risks with nematicides and the need for new approaches. In JA. Veech & D.W. Dickson, eds. *Vistas on nematology*, p. 469-476. Hyattsville, USA, Society of Nematologists.

Keywords: ultra-high dilution, root-knot nematode, *Arabidopsis thaliana*

© International Journal of High Dilution Research.
Not for commercial purposes.



Cite as: Rodiuc N. The effect of ultra-high-diluted drugs on plant-nematode interaction. Proceedings of the XXIX GIRI Meeting; 2015 June 3 – 5; Verona (Italy). *Int J High Dilution Res.* 2015; 14(2): 9-10